arXiv:1605.06777v2 [astro-ph.SR] 3 Jun 2016

Constraints on the presence of SiO gas in the debris disk of HD 172555

T. L. Wilson[1], R. Nilsson[2,3], C. H. Chen[4], C. M. Lisse[5], M. Moerchen[4], H.-U. Käufl[6], A. Banzatti[4]

Received _____; accepted _____

Accepted by ApJ May 3, 2016

[1]United States Naval Research Laboratory, Washington, DC 20375

[2]Astrophysics Department, American Museum of Natural History, Central Park West at 79th Street, New York, NY 10024, USA

[3]Department of Astronomy, Stockholm University, AlbaNova University Center, Roslagstullsbacken 21, 106 91 Stockholm, Sweden

[4]Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21212

[5]Johns Hopkins Applied Physics Laboratory, Silver Spring MD, 20723

[6]European Southern Observatory, K-Schwarzschild-str. 2, Garching, Germany, 85748

ABSTRACT

We have carried out two sets of observations to quantify the properties of SiO gas in the unusual HD 172555 debris disk: (1) a search for the $J = 8 - 7$ rotational transition from the vibrational ground state, carried out with the APEX sub-millimeter telescope and heterodyne receiver at 863 μm, and (2) a search at 8.3 μm for the P(17) ro-vibrational transition of gas phase SiO, carried out with VLT/VISIR with a resolution, $\lambda/\Delta\lambda$, of 30000. The APEX measurement resulted in a 3 σ non-detection of an interstellar feature, but only an upper limit to emission at the radial velocity and linewidth expected from HD 172555. The VLT/VISIR result was also an upper limit. These were used to provide limits for the abundance of gas phase SiO, for a range of temperatures. The upper limit from our APEX detection, assuming an 8000 K primary star photospheric excitation, falls more than an order of magnitude below the self-shielding stability threshold derived by Johnson et al. (2012). Our results thus favor a solid-state origin for the 8.3 μm feature seen in the Spitzer IRS spectrum of the circumstellar excess emission, and the production of circumstellar O I and Si I by SiO UV photolysis. The implications of these estimates are explored in the framework of models of the HD 172555 circumstellar disk.

Subject headings: circumstellar matter – stars: individual (HD 172555) – submillimeter: planetary systems – techniques: spectroscopic

1. Introduction

The occurrence rate of *debris disks* containing hot dust around sun-like stars is very low, ∼2% (Bryden et al. 2006a,b; Chen et al. 2006). Recent work suggests that the hot dust around most mature sun-like stars is transiently regenerated, since it is found to be present in quantities far in excess of that expected to arise from sublimation of comets (Beichman et al. 2005) or slow collisional grinding of asteroids left over from the era of planet formation (Wyatt et al. 2007). Terrestrial planets are thought to grow by accumulation of smaller objects, through the process of collisional accretion (Kenyon & Bromley 2004, 2006) on timescales of 10 to 100 Myr (see Wetherill 1990; Yin et al. 2002; Chambers 2004, and references therein).

As we will discuss, HD 172555, located at a distance of 29.3 pc and belonging to the β Pic Moving Group, is an unusual debris disk system. At an age of more than 15 Myr (Mamajek et al. 2014; Bell & Mamajek 2015), it is at a stage of development where the dusty primordial disk should have dissipated and planets formed. Yet, HD 172555 has been identified as a source with pronounced mid-IR excess emission (Schütz et al. 2005; Chen et al. 2006), which is a signpost of thermally radiating circumstellar dust. Dust and gas present at this late stage of disk evolution is likely secondary, i.e. produced from bodies already aggregated, via impacts and cometary outgassing, forming a debris disk. Collisions of planet-sized objects in debris disks may be distinguished from slow, collisional grinding since the giant collisions produce much larger masses of warm dust than that expected from steady-state collisional evolution (Wyatt et al. 2007, 2011). Unlike most young debris disks, and debris disks in general, the dust surrounding the primary star of HD 172555 is peculiar. Lisse et al. (2009) took Spitzer/IRS spectra with $\lambda/\Delta\lambda = 90$, covering wavelengths λ = 5–35 μm and combined these results with additional IR photometry to determine the Spectral Energy Distribution (SED) from 5 μm to 100 μm. They showed that a large

amount of fine dust with a temperature of 245 K was at 6 AU from the primary, consistent with the later measurements of Pantin & di Folco (2011) using "Lucky Imaging", and Smith et al. (2012) using VLT/MIDI interferometry that showed that the circumstellar dust emission arises from an axisymmetric torus of radius 5.8±0.6 AU (0.″2 at 29.3 pc). In addition, from blended 8 μm features Lisse et al. (2009) reported that 10^{47} SiO molecules were present in the gas phase and proposed that the most likely source of the fine dust and gas phase SiO was a massive hypervelocity impact between two rocky planetoids which liberated huge amounts of melted and reformed ejecta. The dust mineralogy is composed primarily of highly refractory, nonequilibrium materials, with approximately three quarters of the Si atoms in silica (SiO_2) species. Tektite and obsidian laboratory thermal emission spectra (non equilibrium glassy silicas found in impact and magmatic systems) are required to fit the data (Lisse et al. 2009). The best-fit model size distribution for the observed fine dust is a number-size distribution of

$$dn/da = a^{-3.95\pm0.10}$$

While IR photometry of the system has remained stable since the 1983 IRAS mission, this steep size distribution, with abundant micron-sized particles, argues for a fresh source of material within the last 0.1 Myr (Lisse et al. 2009; Johnson et al. 2012).

In an analysis of the HD 172555 data, Johnson et al. (2012) argued that the 5.8 AU torus containing the $\sim 10^{47}$ molecules of gas phase SiO reported by Lisse et al. (2009) would have to be optically thick in the UV to provide self-shielding stability over decades for any circumstellar SiO gas, while being optically thin in the IR, to allow detection by Spitzer but reducing any radiation pressure forces on the dust. Such a torus should also contain an abundance of silicon and oxygen atoms. Johnson et al. (2012) found that about 10^{48} atoms of Si and O vapor would be needed to prevent the photodissociation of SiO. HD 172555 has been re-classified as an A6V–A7V star, which would lower the expected

UV flux, and consequently lower the amount needed for stability by about a factor of two. O I in abundance was detected at 63 μm by Riviere-Marichalar et al. (2012). More recently, Kiefer et al. (2014) reported the detection of time variable Ca II K and H doublet lines, possibly associated with falling evaporating bodies (FEBs or exocomets), which may provide an alternate explanation for the fine dust and gas phase SiO in the HD 172555 system.

To refine models for HD 172555, we have searched for ro-vibrational lines of SiO at 8.3 μm with VLT/VISIR with finer velocity resolution than was possible with Spitzer. [1] The VISIR measurements are most sensitive to gas with effective ro-vibrational temperatures up to 500 K, causing the flux to concentrate in a few lines near the ground state. It is also possible that the gas phase SiO might be as cold as the dust temperature, ~ 245 K. To investigate this possibility, we also searched for the $J = 8 - 7$ rotational transition with the 12-m APEX submillimeter telescope.[2]

2. Observations and Data Reduction

2.1. APEX

Data were obtained during August, November and December 2014 using the 12-meter Atacama Pathfinder EXperiment (APEX) radio telescope with the APEX-2 heterodyne

[1] This publication is partially based on data acquired with observations collected at the European Organization for Astronomical Research in the Southern Hemisphere, Chile (087.C-0669(B)).

[2]This publication is partially based on data acquired with the Atacama Pathfinder Experiment (APEX). APEX is a collaboration between the Max-Planck-Institut fur Radioastronomie, the European Southern Observatory, and the Onsala Space Observatory.

receiver, which operates in the 267–378 GHz band. There are two SiO transitions from the vibrational ground state, at 303.93 GHz ($J = 7 - 6$) and at 347.33 GHz ($J = 8 - 7$). We chose the $J = 8 - 7$ transition since this will provide a larger signal for a given column density of SiO gas warmer than 50 K. There are thousands of lines from exotic molecular species within $\pm 20\,\mathrm{km\,s^{-1}}$ of the center of the spectrometer band (from a perusal of the Splatalogue website)[3], but all are rather unlikely to be present. The $J = 8 - 7$ rotational transition from the vibrational ground state of SiO has a rest wavelength of 863 μm or frequency of $\nu = 347.33$ GHz. At that frequency, the full width to half power (FWHP) beam size of the instrument was $17''$, with a velocity resolution of $1.3\,\mathrm{km\,s^{-1}}$. The spectrometer was centered on $1\,\mathrm{km\,s^{-1}}$, the velocity of the star in the Local Standard of Rest (LSR) frame. The receiver noise was less than 200 K and the system noise, including the Earth's atmosphere was 350 K during the measurements. The integration time was 4.8 hours, on-source, with an equal amount of off-source integration time in the position-switching mode, with an off-source position $10'$ from the source position. Calibration was made using the chopper wheel procedure (see, e. g. Wilson et al. 2013). The GILDAS software package with CLASS commands was used for reducing and adding data from all scans, performing baseline subtraction, and fitting a Gaussian to the flux at the line position.[4]

2.2. VLT

The P(17) transition from the ($v = 1$, $J = 17$) to the ($v = 0$, $J = 18$) energy level of SiO gas lies in the mid-IR with a rest wavelength of $\lambda = 8.305$ μm, or $\nu = 3.61 \times 10^{13}$ Hz. Usable VLT/VISIR data were taken on August 19 and 20, 2010, from Paranal in fair

[3]See `http://www.cv.nrao.edu/php/splat/`

[4]See `http://www.iram.fr/IRAMFR/GILDAS`.

weather.

The VISIR spectrometer was used in the high resolution cross-dispersed chopping mode, with a resolution of $\lambda/\Delta\lambda = 3 \times 10^4$, so each channel has a width of $10\,\mathrm{km\,s^{-1}}$. At 8.3 μm, the diffraction limited Full Width at Half Power (FWHP) beam size is 0.″2. The total integration time was 3220 s. The bandpass correction for telluric lines was determined from measurements of α Cen. The offset from zero level in the VLT data from chopping was set equal to the flux density from the broadband Spitzer spectrum for this wavelength, 1.47 Jy, to establish the calibration (see Fig. 1 of Johnson et al. 2012). The VISIR data were reduced using a suite of IDL routines built to optimize the standard VISIR data reduction of the ESO Recipe Execution Tool (EsoRex, version 3.9.0), as described in Banzatti et al. (2014). The final spectrum is shown in Fig. 2, and the results are discussed further in Section 3.

The line parameters are listed in Table 1. Our estimates of the number of SiO molecules in the gas phase are derived in Section 3 below.

3. Results

3.1. The APEX Data

From Wilson et al. (2013), the relation for a two level system is:

$$N_{\rm l} = 93.5 \frac{g_{\rm l}\,\nu^3}{g_{\rm u}\,A_{\rm ul}} \frac{1}{[1 - e^{(-4.80\times10^{-2}\nu/T_{\rm ex})}]} \int \tau \mathrm{d}V, \qquad (1)$$

where $N_{\rm l}$ is the beam averaged column density in the lower level, ν is the line frequency in GHz, $g_{\rm l}$ is the statistical weight of the lower level, $g_{\rm u}$ of the upper level, $\mathrm{A_{ul}}$ is the Einstein coefficient, τ is the apparent optical depth and V is the velocity width in $\mathrm{km\,s^{-1}}$. The

excitation temperature, T_{ex}, characterizes the populations of the upper and lower levels:

$$\frac{n_u}{n_l} = \frac{g_u}{g_l} e^{(-E/T_{ex})}, \tag{2}$$

where n_u is the population of the upper level, n_l the population of the lower level, and E is the energy of n_u above n_l in K. For the 863 µm line, we can simplify Eq. 1 with the use of $h\nu \ll kT$:

$$N_l = 1.95 \times 10^3 \frac{g_l \nu^2}{g_u A_{ul}} \int \tau dV, \tag{3}$$

Further, we have $T_{MB} = T_{ex}\tau$, $A_{ul} = 2.2 \times 10^{-3}\,s^{-1}$, g_l=15, g_u=17, and $\nu = 347.333$ GHz. Then the column density in the $J = 7$ level is given by:

$$N_7 = 1.0 \cdot 10^{11} \cdot T_{MB}\,\Delta V \tag{4}$$

The parameters are given in Table 1. A radial velocity of -22 $km\,s^{-1}$ is consistent with the interstellar lines of sodium and calcium found by Kiefer et al. (2014) when observing in the direction of HD 172555. With the detection of a single line, however, no detailed analysis is possible on this potential ISM source.

In the following, we restrict our analysis to the SiO emission assumed to arise from circumstellar material in the HD172555 system. This should have a V_{LSR}=1 $km\,s^{-1}$ and a FWHP of 33 $km\,s^{-1}$ (Gontcharov 2006; Kiefer et al. 2014). Qualitatively, from a perusal of Fig. 1, no line seems to be present above the noise at this relative velocity. Quantitatively, we employed a Gaussian fit, using V_{LSR}=1 $km\,s^{-1}$ and a FWHP of 33 $km\,s^{-1}$ and find an upper limit of T_A^*=2.3mK, with an RMS spectral noise of T_A^*=4.63mK. This is an underestimate of the upper limit to any signal, so we have used the 3 RMS noise limit, T_A^*=13.9mK in the following. From this limit to T_A^* and the assumed linewidth, we determined the number of SiO molecules in the $J = 7$ rotational level of the vibrational ground state. This is

$$N_7 \leq 5.5 \times 10^{10}\,cm^{-2} \tag{5}$$

In the following equations we use equal signs, although this is an upper limit. The total number of SiO molecules in all rotational and vibrational levels depends on temperature. This number is estimated using the following analysis. The total column density in the ground vibrational state is related to the total column density in all rotational levels in the vibrational ground state, $N(\mathrm{total},\, v=0)$, by:

$$N(\mathrm{total}, v=0) = \frac{N_J}{g_J} \times \mathrm{e}^{(E_J/T_{\mathrm{rot}})} \times \left(\sum_J g_J \, \mathrm{e}^{E_{\mathrm{rot}}/T} \right), \tag{6}$$

where B_e=1.05 K (the molecular constants of SiO are to be found in Barton et al. (2013). Here these are expressed in temperature units. The sum can be replaced by an integral for E_{rot}/k greater than 5 K. [5] Then for $J = 7$ we have:

$$N(\mathrm{total}, v=0) = \frac{N_7}{15} \times \left(\frac{T}{1.05} \right) \times \mathrm{e}^{(43.6/T)}, \tag{7}$$

The total column density is the sum over the populations of the vibrational states, given by:

$$N(\mathrm{total}, \mathrm{vibrational}) = \sum_{v=0} \mathrm{e}^{E_{\mathrm{vib}}/T} \tag{8}$$

where $E_{vib} = 1758$ K is the energy of the $v = 1$ state above the ground state, in K, and the vibrational levels are very nearly equally spaced by this energy. The column density of SiO molecules, in cm^{-2}, is the product of Eq. 5, Eq. 7, and Eq. 8. The total number of SiO molecules is the product of the column density and and the beam area in cm^2. The FWHP beam of APEX at the line frequency is 17.″6; taking this as the diameter, the beam area is $4.7 \times 10^{31}\,\mathrm{cm}^2$. The upper limit to the total number of SiO molecules is plotted as a function of temperature in Fig. 3. We allow a wide possible range of excitation temperatures, since arguments have been made in the literature by Lisse et al. (2009) for the excitation of SiO at 8000 K, the stellar surface temperature, and by Johnson et al. (2012) for an excitation at 245 K by intermixed dust. For two extreme values, 300 K and 8000 K, the 3 σ limit for the total number of SiO molecules is $\leq 5.5 \times 10^{43}$ and $\leq 4.2 \times 10^{45}$, respectively.

[5] see http://www.exomol.com/exomol2/data/SiO/28Si-16O/EBJT/28Si-16O

3.2. The VLT Data

There is no obvious spectral line in the 8.3 μm VLT/VISIR spectrum, so we have used three times the RMS limit in a single spectral channel as the limit. This is 0.48 Jy in a 10 km s^{-1} channel. For the P(17) transition, the wavelength λ is 8.308 μm. For the following analysis, we follow the approach used in Lisse et al. (2009). This can be obtained from Eq. 1 if $h\nu \ll kT_{\rm ex}$. Using $\mathfrak{J}$, the spontaneous emission coefficient in the equation of radiative transfer, the Einstein A coefficient between the upper level, $J = 18$, $v = 1$, and lower level, $J = 17$, $v = 0$:

$$\mathfrak{J}_u = \frac{h\nu}{4\pi} A_{\rm ul} N \tag{9}$$

where N is the column density in units of cm^{-2} and h is Planck's constant.

$$F = \int S_\nu {\rm d}\nu = \frac{h\nu}{4\pi} A_{\rm ul} N \Delta\Omega \tag{10}$$

where N is the column density in the upper level (ul). The total number of molecules, is estimated from the fraction in the observed level. For this, the summation (Eq. 6) cannot be replaced by an integral. For the VLT, the area subtended by the FWHP of the diffraction beam is 6×10^{27} cm^{-2} for a distance of 29.3 pc. The 3 σ limit for the flux density in the 8.3 μ line is $\leq$0.48 Jy, in a 10 km s^{-1} wide channel. The other parameters are $v = 1$, $J_{\rm u} = 17$, $J_{\rm l} = 18$, and $A_{\rm ul} = 3.44\,s^{-1}$, from the Barton et al. (2013) molecular data.

Estimates of the total number of SiO molecules in the gas phase, from our data and from Lisse et al. (2009), are plotted in Fig. 3 as a function of temperature. Our APEX data in Fig. 1 was analyzed using Eq. 5, Eq. 6 and Eq. 8, under the assumption of LTE. For $T = 8000$ K, the curve based on the APEX data is more than a factor of more than 40 below the value reported by Lisse et al. (2009), while the VLT limit is a factor of more than a factor of 10^3 below the value of Lisse et al. (2009). For $T = 300$ K, the APEX data is more significant, being more than a factor of 10^3 below the value of Lisse et al. (2009).

4. Discussion

The 2.3 km s^{-1}wide line at V_{LSR}=-22 km s^{-1} arises in the interstellar medium (see, e.g., Kiefer et al. 2014), so is not related to gas in the disk associated with HD 172555. Although weak, the radial velocity is in agreement with the value for the interstellar atomic lines found by Kiefer et al. (2014). We will not discuss this feature in the following, but note that this result shows that the APEX system has functioned satisfactorally during our observations. In the following, we concentrate on the upper limit for SiO at V_{LSR}=0 km s^{-1}.

According to Johnson et al. (2012), the abundance must exceed 10^{47} to allow survival against photo dissociation of the SiO in the environment of HD 172555. With the re-assessment of the classification of the central star, the amount of SiO is lower, but the VLT data sets an upper limit much lower than 10^{46} for T $\geq$ 300 K while the APEX limit is below this value for all temperatures considered. Taken together, these results indicate that the abundance based on the feature in the Spitzer spectrum of Lisse et al. (2009) should be reduced. On the basis of the gas disk analysis of Johnson et al. (2012), it is likely that SiO in the gas phase would not survive; instead any SiO gas produced by a giant hypervelocity impact should be photolyzed into Si I + O I atoms by the strong UV flux of the A7V primary, producing the O I population detected by Herschel (Riviere-Marichalar et al. 2012). To explain the 8 μm emission feature seen in the Spitzer data, Johnson et al. (2012) suggested an alternative explanation for its source: this would be a shoulder seen from thermal emission produced by fine grained, amorphous, hydrated silica (i.e., opalescent) smokes, as seen in laboratory spectra by Speck et al. (2011) and as detected on the surface of Mars by Ruff et al. (2011) and Sun et al. (2015).

If the abundance of gas phase SiO is close to our limit, there are a number of additional findings. First, the dissociated SiO gas could be a source of the O I detected by Herschel, so we do not require substantial CO or CO_2 gas to do so. The lifetime against photodestruction

for O I is considerably longer than that of gas phase SiO, so the abundance of gas phase SiO would have been larger in the past, so the presence of O I is consistent with a hypervelocity collision 10^4 years ago (Johnson et al. 2012). Alternatively, Riviere-Marichalar et al. (2012) speculate that the O I may have been produced by the slow destruction of grains, that is, collisions, sublimation or sputtering, and remained in place since this species is less affected by radiation pressure than SiO. Second, the heating and vaporization of rocky dust material that produces SiO could also produce the Ca II and Na I detected by Kiefer et al. (2014), since a process that destroys the SiO lattice of silicate dust should also liberate the associated cations in the matrix.

Our abundance is based on the 3σ upper limits from two non-detections. A reasonable follow-up to go deeper and search for a detection of total SiO abundances down to 10^{42}, while also mapping the system in the dust continuum, could be provided by a short integration (1 hr, including calibration) with the ALMA array using a resolution of $0.2''$. SiO abundances in the 10^{42} range are expected if the Keifer et al. (2014) detection of Ca II and Na I are due to the destruction of rocky material in-system.

5. Conclusions

We have a 3.3σ detection of the $J = 8 - 7$ rotational line (Fig. 1) from the interstellar medium toward HD 172555. For gas associated with HD 172555 itself, we have the 3σ limits for the $J = 8 - 7$ rotational line and the P(17) ro-vibration line at 8.3 μm of gas phase SiO, the conclusions are:

1. From the APEX data, our upper limit for 300 K, the limit is 5.3×10^{43}. The APEX data is more sensitive to cooler gas, so these results preclude large amounts of cooler gas phase SiO. For $T = 8000$ K, the total LTE abundance of gas phase SiO is less

than 4.4×10^{45}.

2. From the VLT data our upper limit for the total LTE is less than 2.7×10^{47} at 8000 K. At lower temperatures, the abundance is smaller. See Fig. 3 for the dependence of abundance on temperature.

3. The upper limit obtained with APEX, if interpreted as representative of the minimum past SiO gas content of the system, does not exclude SiO as a source of the O I detected with Herschel, so substantial amounts of CO or CO_2 gas are not required.

4. The heating and vaporization of rocky dust material that could produce gas phase SiO to the upper limit found with APEX could also produce the Ca II and Na I detected by Kiefer et al. (2014).

5. With the SiO gas stability analysis of Johnson et al. (2012), revised down to about 5×10^{48} molecules for the approximately two times lower UV flux for an A6V type primary star, our upper limit on SiO gas suggests that the feature in the Spitzer spectrum previously attributed to gas phase SiO is caused by an amphorous silica compound.

We thank an anonymous referee for helpful comments on the text and interpretation. R.N. was funded by the Swedish Research Council's International Postdoctoral Grant No. 637-2013-474.

APEX (APEX-2), VLT (VISIR).

REFERENCES

Banzatti, A., Meyer, M. R., Manara, C. F., Pontoppidan, K. M., Test, L. 2014 ApJ, 780, 26

Barton, E., Yunchenko, S. N. & Tennyson, J. 2013, MNRAS, 434, 1469

Beichman, C. A., Bryden, G., Gautier, T., et al. 2005 ApJ, 626, 1061

Bell C. P. M. & Mamajek, E. E. 2015, MNRAS, 454, 593

Bryden, G., Beichman, C. A., Trilling, D. E., et al. 2006a ApJ, 636, 1098

Bryden, G., Beichman, C. A., Carpenter, J. M., et al. 2006b ApJ, 705, 1226

Chambers, J.E. 2004 Earth & Planetary Sci Lett 223, 241

Chen, C. H., Sargent, B. A., Bohac, C., et al. 2006, ApJS, 166, 351

Dumke, M. & Mac-Auliffe, F. 2010, Proc of SPIE, 7737, 77371J-2

Gontcharov, G. A. 2006 Astronom Lett 32, 759

Johnson, B. C., Lisse, C. M., Chen, C. H., et al. 2012, ApJ 761, 45

Kenyon, S. J. & Bromley, B. C. 2004 ApJ, 602, L133

Kenyon, S. J. & Bromley, B. C. 2006 AJ, 131, 1837

Kiefer, F., Lecavelier des Etangs, A., Augereau, J.-C., et al. 2014, A&A, 561, L10.

Lisse, C. M., Beichman, C. A., Bryden, G. & Wyatt, M. C. 2007, ApJ, 658, 584

Lisse, C. M., Chen, C. H., Wyatt, M. C., 2009, ApJ, 701, 2019

Mamajek, E. E., Bell C. P. M. & Cameron, P. M. 2014, MNRAS, 445, 2169

Moór, A., Ábrahám, P., Juhász, A. et al. 2011, ApJ, 740, L7

Pantin, E. & di Folco, E. 2011, EPSC-DPS Joint Meeting 2011 Nantes, France

Riviere-Marichalar, P., Barrado, D., Augereau, J.-C., et al. 2012, A&A, 546, L8

Ruff, S. W., Farmer, J. D., Calvin, W. M., et al., 2011 J. Geophys. Res., 116, E00F23

Schütz, O., Meeus, G., & Sterzik, M. F. 2005, A&A, 431, 175

Smith, R., Wyatt, M. C., & Haniff, C. A. 2012, MNRAS, 422, 2560

Speck, A., Whittington, A. G. & Hofmeister, A. M. 2011, ApJ, 740, 93

Sun, V. Z., Milliken, R. E., Rieff, S. W. & Farmer, J. D. 2015, Lunar & Planet. Sci. Conf. 46, 2157

Wetherill, G. W. 1990 Annu. Rev. Earth Planet. Sci. 18, 205

Wilson, T. L., Rohlfs, K. & Huettemeister, S. 2013, Tools of Radio Astronomy, Springer, Berlin

Wyatt, M. C., Smith, R., Su, K. Y. L. et al. 2007, ApJ, 663, 365

Wyatt, M. C., Clarke, C. J., & Booth, M. 2011, Celestial Mechanics and Dynamical Astronomy, 111, 1

Yin, Q., Jacobsen, S. B., Yamashita, K., Blichert-Toft, J. & Telouk, P. 2002 Nature 418, 949

This manuscript was prepared with the AAS LaTeX macros v5.2.



Fig. 1.— The spectrum taken with the APEX millimeter/sub-mm telescope. The fit in green shows the 3.3 σ feature which is taken as a detection of the SiO $J = 8-7$ rotational transition from the interstellar medium, at V_{LSR}=-22 km s^{-1}. The the fit in red was constrained to be centered at V_{LSR}=1 km s^{-1} and to have a width of 33 km s^{-1} This spectrum is on a T_A^* scale. The peak $T_A^* \leq 2.3$ mill K. To convert to main-beam brightness temperatures, multiply by 1.3. For the estimate of the total number of SiO molecules, a T_A^* equal to three times RMS, 13.9 milli K, was used. The radial velocities are with respect to the Local Standard of Rest using the Standard Solar Motion.



Fig. 2.— The spectrum of the P(17) ro-vibrational line of gas phase SiO, taken with VLT/VISIR with a resolution of 3×10^4. The central vertical line at 8.308 μm shows the expected location of the P(17) line of SiO.



Fig. 3.— A plot of limits for the total abundance of gas phase SiO in HD172555 as a function of the LTE temperature. For T≥500 K, the lowest curve is based on our VLT measurement of the ro-vibrational P(17) line at 8.3 μm. The next higher curve is based on our APEX measurement of the J=8-7 rotational transition at 347.33 GHz. The downward arrows indicated that these values are upper limits. The horizontal line shows the abundance of gas phase SiO reported by Lisse et al. 2009, from their Spitzer data.

Table 1. Results

Telescope	Transition	Wavelength (μm)	S_ν (Jy)	$T_{\rm MB}$ (K)	$\Delta V_{1/2}$ (km s^{-1})	$\Delta V_{\rm LSR}$ (km s^{-1})
(1)	(2)	(3)	(4)	(5)	(6)	(7)
VLT	P(17)	8.3	$\leq 0.48^a$	···	10^b	0^c
APEX	$J = 8 - 7$	863	···	0.018±0.006 K	2.63^d	-22^e
APEX	$J = 8 - 7$	863	···	$\leq 0.017^a$	33^f	0^c

[a] 3σ limit

[b] Velocity resolution

[c] From gas associated with HD 172555

[d] Measured line width

[e] From interstellar gas

[f] Assumed line width